

## Gel

- Individual treatment of digital dermatitis
- No antibiotics
- Strong adhesion to the hoof
- Safe for the user
- No withdrawal times



# Gel

## Hoof problems cause significant economic losses.

Lameness is often perceived as costly when an animal is physically lame and suffering, but in reality, the impact extends far beyond the discomfort of that one animal. For example, in dairy cows, longer calving intervals and mastitis prevalence are often related to poor hoof health conditions and indirectly decrease the financial bottom line. Hoof problems also cause extra labor, high culling and increased treatment costs.

**Lameness costs can easily run up to \$ 400 annually per dairy cow.**

**Hoof-fit Gel is a proven cure against Digital Dermatitis.**

The efficacy of **Hoof-fit Gel** has been independently confirmed in 2 trials :

A European trial on 172 cows showed a 92% cure rate for painful ulcerative stage (M2) lesions of bovine digital dermatitis (BDD). A Canadian trial involving 50 cows showed a near 100% cure rate of M2 lesions.

**Cows with lameness issues need your special care and attention.**

If you are unsure of the treatment required, contact a veterinarian or professional hoof trimmer to treat lame cows before implementing a spraying and/or foot bathing protocol. We recommend using follow up protocols after lame cows have been treated, to optimize hoof health management and achieve best results.

### Usage tips :

- Use the Gel above freezing temperatures for easy brushing.
- Close the Gel container after use.
- Make sure wraps are removed within 3-4 days after treatment.
- Make sure that no foot baths are used until the wraps are removed.



## How to use Hoof-fit Gel



1. Trim the claws 2 to 3 times per year. All the animals should preferably be trimmed by a professional trimmer.



2. Clean the claw thoroughly. Make sure that any manure present is removed from the claw and from the inter-digital space. Dry the claw.



3. Apply **Hoof-fit Gel** to the claw and the inter-digital space using the brush provided.



4. We advise you to use wraps. Make sure that a wrap is removed within 3-4 days.



5. If necessary, apply a thin layer of Gel without a wrap on days 3 and 7.

### Ingredients :

- Activated Copper and Zinc chelate
- Isopropanol

### Hoof fit Gel is available in

- 330 ml jar

DIN : 02441284  
Health Canada